## Price Directed Distributed Optimization and Primal Recovery

Tetiana Parshakova

Yicheng Bai Garrett van Ryzin Stephen Boyd

Amazon SCOT

**INFORMS** 

10/21/24

Price directed distributed optimization

Primal recovery

Numerical results

Appendix

#### Our focus

#### our setting

- substantial computational cost per agent
- small number of iterations is desirable
- care about feasibility, some suboptimality is tolerable

#### we propose (postprocessing) method that

- uses low-precision optimal dual variable
- recovers close to feasible primal point
- uses only parallel calls to agents; avoids sequential calls

## **Distributed optimization**

minimize 
$$f(x)$$
  
subject to  $Ax \le b$ 

- $ightharpoonup x = (x_1, \dots, x_M) \in \mathbf{R}^n$  is variable,  $x_i \in \mathbf{R}^{n_i}$
- $f(x) = \sum_{i=1}^{M} f_i(x_i)$  is block separable
- $lackbox{}{} f_i: \mathbf{R}^{n_i} 
  ightarrow \mathbf{R} \cup \{\infty\}$  convex, closed and proper
- ightharpoonup infinite values of f encode constraints
- $lackbox{ } A=(A_1,\ldots,A_M)\in \mathbf{R}^{m imes n} \ \ ext{and} \ \ b\in \mathbf{R}^m \ \ ext{are given}$

# Conjugate subgradient oracle

▶ for  $y \in \mathbf{R}^m$  (dom  $f^* = \mathbf{R}^m$ ), oracle returns  $x(y) \in \partial f^*(y)$ 

$$-f^*(y) = \inf_{x \in \mathbf{dom}\, f} \left( f(x) - y^T x \right)$$

- lacktriangle no access to function values f(x) or subgradient in  $\partial f(x)$
- neutral cutting plane for dual variable

$$\left\{ \tilde{\lambda} \in \mathbf{R}^m \mid (-Ax(y) + b)^T (\tilde{\lambda} - \lambda) \le 0, \ y = -A^T \lambda \right\}$$

## **Dual problem**

#### solve the dual problem

- ▶ subgradient methods (Shor, 1962)
- ► localization methods
  - ► analytic center cutting-plane method (ACCPM)
  - maximum volume ellipsoid cutting-plane method
  - Chebyshev center cutting-plane method
- ▶ we've settled on proximal point ACCPM

# **Optimality conditions**

KKT conditions

$$x \in \partial f^*(y), \ y = -A^T \lambda \tag{1}$$

$$Ax \le b \tag{2}$$

$$\lambda_j (Ax - b)_j = 0, \quad j = 1, \dots, m \tag{3}$$

$$\lambda \ge 0 \tag{4}$$

lacktriangle primal and complementary slackness residuals for a pair  $(x,\lambda)\in \mathbf{R}^n imes \mathbf{R}^m_+$ 

$$r_p = \mathbf{1}^T (Ax - b)_+, \qquad r_c = \lambda^T |Ax - b|$$

Price directed distributed optimization

Primal recovery

Numerical results

**Appendix** 

Primal recovery 8

# **Primal recovery**

given current prices  $y^k = -A^T \lambda^k = (y_1^k, \dots, y_M^k)$ 

- 1. agent i returns multiple offers  $z_i^{(1)},\dots,z_i^{(N)}$  wrt  $y_i^k=-A_i^T\lambda^k$ 
  - $-f_i^*(y_i^k) \le f_i(z_i) y_i^{kT} z_i \approx -f_i^*(y_i^k)$
  - computed in parallel (clock time equivalent to single response)
- 2. central node constructs a convex combination of offers  $\bar{x}$  minimizing the residuals

minimize 
$$\begin{array}{ll} r_p + r_c \\ \text{subject to} & \bar{x}_i = Z_i u_i, \quad i = 1, \dots, M \\ & \mathbf{1}^T u_i = 1, \ u_i \geq 0, \quad i = 1, \dots, M \\ & \bar{x} = (\bar{x}_1, \dots, \bar{x}_M) \end{array}$$

don't need to run primal recovery at every step k (periodic runs sufficient)

# Approximate conjugate subgradient oracle

• for any y, an  $\epsilon_v$ -value suboptimal primal variable  $x^{\mathsf{v}}(y)$ 

$$-f^*(y) \le f(x^{\mathsf{v}}(y)) - y^T x^{\mathsf{v}}(y) \le -f^*(y) + \epsilon_v |f^*(y)|$$

- ▶ any convex combination  $\bar{x}$ :  $f(\bar{x}) y^T \bar{x} \leq -f^*(y) + \epsilon_v |f^*(y)|$
- for any y, primal variable  $x^{p}(y)$  with  $\epsilon_{p}$ -perturbed prices

$$f(x^{\mathsf{p}}(y)) - (y+\delta)^T x^{\mathsf{p}}(y) = -f^*(y+\delta), \quad \delta \in [-\epsilon_p |y|, \epsilon_p |y|]$$

• for L-Lipschitz  $f^*$ , for any convex combination  $\bar{x}$ :  $f(\bar{x}) - y^T \bar{x} \leq -f^*(y) + \epsilon_p L \|y\|$ 

Primal recovery 10

Price directed distributed optimization

Primal recovery

Numerical results


Appendix


## Setting

- $\blacktriangleright$  run localization method for solving the dual, compute  $\lambda^k$
- lacktriangle price directed interface gives  $x^k \in \partial f^*(-A^T\lambda^k)$
- ightharpoonup  $ar{x}$  returned by primal recovery method
- ightharpoonup N=10 suboptimal offers with  $\epsilon=10\%$ -suboptimality
- ightharpoonup primal recovery effective when  $m \ll n$

## **Resource allocation**


- ightharpoonup M=100 agents, m=8 resources, n=800
- $ightharpoonup \epsilon_v = 10\%$ -value suboptimality






#### Resource allocation


- ightharpoonup M=100 agents, m=8 resources, n=800
- $ightharpoonup \epsilon_p = 10\%$ -perturbed prices






# **Assignment problem**


- ightharpoonup M = 208 agents, m = 8, n = 1608
- $ightharpoonup \epsilon_v = 10\%$ -value suboptimality






# Assignment problem


- ightharpoonup M=208 agents, m=8, n=1608
- $ightharpoonup \epsilon_p = 10\%$ -perturbed prices






# **Linear programming**


- ightharpoonup M = 100 agents, m = 8, n = 800
- $ightharpoonup \epsilon_v = 10\%$ -value suboptimality






# **Linear programming**

- ightharpoonup M = 100 agents, m = 8, n = 800
- $ightharpoonup \epsilon_p = 10\%$ -perturbed prices





Price directed distributed optimization

Primal recovery

Numerical results

Appendix

Appendix 19

## Primal recovery method

lower bound

$$\mathcal{L}(x^k, \lambda^k) = \operatorname*{argmin}_x (f(x) + \lambda^{kT} (Ax - b)) = -f^*(-A^T \lambda^k) - \lambda^{kT} b \leq p^*$$

ightharpoonup  $\epsilon_v$ -value suboptimality

$$\mathcal{L}(\bar{x}^{\mathsf{v}}, \lambda^k) = f(\bar{x}^{\mathsf{v}}) + \lambda^{kT} (A\bar{x}^{\mathsf{v}} - b) \le p^{\star} + \epsilon_v \sum_{i=1}^{M} |f_i^{\star}(-A_i^T \lambda^k)|$$

ightharpoonup  $\epsilon_p$ -price perturbation

$$\mathcal{L}(\bar{x}^{\mathsf{p}}, \lambda^k) = f(\bar{x}^{\mathsf{p}}) + \lambda^{kT} (A\bar{x}^{\mathsf{p}} - b) \le p^{\star} + \epsilon_p L \sum_{i=1}^{M} \|A_i^T \lambda^k\|$$

Appendix 20